

ORIGINAL RESEARCH

Comparing clinical presentation, viremia, and immunological factors at various severity presentations in hospitalized children affected by COVID-19: A cross-sectional study

Marzieh Jamalidoust¹  | Seyedeh Sedigheh Hamzavi^{2,3}  | Eslam Shorafa⁴  |
Mandana Namayandeh¹  | Laiba Batool³ | Seyedeh Narges Abootalebi^{4,5} 

¹Department of Virology, Professor Alborzi Clinical Microbiology Research Center, Namazi Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

²Professor Alborzi Clinical Microbiology Research Center, Namazi Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

³School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

⁴Pediatric Intensivist, Intensive Care Unit division, Department of Pediatrics, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

⁵Biotechnology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Correspondence

Marzieh Jamalidoust, Department of Virology, Alborzi Clinical Microbiology Research Center, Namazi Hospital, Shiraz University of Medical Sciences, P.O. Box 31, Shiraz 71937-11351, Iran.

Email: mjamalidoust@gmail.com

Seyedeh Sedigheh Hamzavi, Alborzi Clinical Microbiology Research Center, Namazi Hospital, Shiraz University of Medical Sciences, P.O. Box 31, Shiraz 71937-11351, Iran.

Email: S.hamzavi55@yahoo.com

Funding information

Shiraz University of Medical Sciences, Clinical Microbiology Research Center

Abstract

Background and Aims: Although SARS-CoV-2 infection usually leads to mild COVID-19 in children, sometimes it causes serious complications, especially in those with underlying diseases. Several factors have been identified in determining disease severity in adults, and limited studies have been conducted in children. The prognostic implications of SARS-CoV-2 RNAemia as an important factor in determining disease severity in children are not well understood.

Methods: In this study, we aimed to prospectively assess the relationship between disease severity and immunological factors and viremia in 47 COVID-19 hospitalized children. In this research, 76.5% of children experienced mild and moderate COVID-19, while 23.5% experienced severe and critical forms of the disease.

Results: The presence of underlying diseases in different groups of pediatric patients differed significantly from each other. On the other hand, clinical symptoms such as vomiting and chest pain as well as laboratory parameters including erythrocyte sedimentation rate were significantly different in different groups of patients. Viremia was seen in only two children, and this had no significant relationship with the severity of COVID-19.

Conclusion: In conclusion, our data confirmed that COVID-19 severity differed in SARS-CoV-2 infected children. Some clinical presentation and lab data parameters were different in various presentation of patients. Viremia was not associated with severity in our study.

KEYWORDS

COVID-19, paraclinical data, pediatrics, SARS-CoV-2, severity, viremia

patients. Our results suggest that ESR is the only factor that distinguishes COVID-19 disease severity among immunological factors and, in contrast to many studies in adult, no significant association was observed between viremia and disease severity in children.

AUTHOR CONTRIBUTIONS

Marzieh Jamalidoust: Conceptualization; project administration; software; supervision; visualization; writing—original draft; writing—review and editing. **Seyedeh Sedigheh Hamzavi:** Conceptualization; funding acquisition. **Eslam Shorafa:** Conceptualization; investigation; supervision. **Mandana Namayandeh:** Data curation; methodology. **Laiba Batool:** Data curation; visualization. **Seyedeh Narges Abootalebi:** Data curation; investigation.

ACKNOWLEDGMENTS

The authors would like to thank Shiraz University of Medical Sciences, Shiraz, Iran and also the Center for Development of Clinical Research of Nemazee Hospital and Dr. Amir Yousef Farahmandi for editorial assistance and Miss. Farhadi for statistical analysis. The study is financially supported by the Shiraz University of Medical Sciences.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ETHICS STATEMENT

The project has been approved by Shiraz University of Medical Sciences (Ethics code Number: IR.SUMS.REC.1400.691). All authors acquiesced to the final version of the manuscript.

TRANSPARENCY STATEMENT

The lead author Marzieh Jamalidoust, Seyedeh Sedigheh Hamzavi affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

ORCID

Marzieh Jamalidoust  <http://orcid.org/0000-0002-7034-1236>

Seyedeh Sedigheh Hamzavi  <http://orcid.org/0000-0002-7081-0127>

Eslam Shorafa  <http://orcid.org/0000-0002-8416-586X>

Mandana Namayandeh  <http://orcid.org/0000-0003-1947-1024>

Seyedeh Narges Abootalebi  <http://orcid.org/0000-0002-1688-4897>

REFERENCES

- Aliabadi N, Marandi NH, Jamalidoust M, et al. Seroprevalence of anti-SARS-CoV-2 antibodies in high-risk occupational and low-risk groups in southwestern Iran. *Jundishapur J Microbiol.* 2022;15(7):e126975.
- Jalil M, Ashkan Z, Gholamnezhad M, et al. Effect of COVID-19 on healthcare workers' morbidity and mortality compared to the general population in Kohgiluyeh and Boyer-Ahmad Province, Iran. *Health Sci Rep.* 2023;6(1):e961.
- Prebensen C, Myhre PL, Jonassen C, et al. SARS-CoV-2 RNA in plasma is associated with ICU admission and mortality in patients hospitalized with COVID-19. *Clin Infect Dis.* 2020;73:e799-e802.
- Takács AT, Bukva M, Gavallér G, et al. Epidemiology and clinical features of SARS-CoV-2 infection in hospitalized children across four waves in Hungary: a retrospective, comparative study from March 2020 to December 2021. *Health Sci Rep.* 2022;5(6):e937.
- Jin Y, Yang H, Ji W, et al. Virology, epidemiology, pathogenesis, and control of COVID-19. *Viruses.* 2020;12(4):372.
- Tsabori S, Makis A, Kosmeri C, Siomou E. Risk factors for severity in children with coronavirus disease 2019: a comprehensive literature review. *Pediatr Clin North Am.* 2021;68(1):321-338.
- Bermejo-Martin JF, González-Rivera M, Almansa R, et al. Viral RNA load in plasma is associated with critical illness and a dysregulated host response in COVID-19. *Crit Care.* 2020;24(1):691.
- Hogan CA, Stevens BA, Sahoo MK, et al. High frequency of SARS-CoV-2 RNAemia and association with severe disease. *Clin Infect Dis.* 2021;72(9):e291-e295.
- Jacobs JL, Mellors JWCID. *Detection of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) RNA in Blood of Patients with Coronavirus Disease 2019 (COVID-19): What Does it Mean?*. Oxford University Press; 2021:e2898-e2900.
- Veyer D, Kernéis S, Poulet G, et al. Highly sensitive quantification of plasma SARS-CoV-2 RNA sheds light on its potential clinical value. *Clin Infect Dis.* 2020;73(9):e2890-e2897.
- Xu D, Zhou F, Sun W, et al. Relationship between serum SARS-CoV-2 nucleic acid (RNAemia) and organ damage in COVID-19 patients: a cohort study. *Clin Infect Dis.* 2020;73(1):68-75.
- Fajnzylber J, Regan J, Coxen K, et al. SARS-CoV-2 viral load is associated with increased disease severity and mortality. *Nat Commun.* 2020;11(1):1-9.
- Andersson MI, Arancibia-Carcamo CV, Auckland K. SARS-CoV-2 RNA detected in blood products from patients with COVID-19 is not associated with infectious virus. *Wellcome Open Res.* 2020;5:181.
- Ghotbabadi SH, Mollaie M, Hamzavi SS, Dashti AS. Clinical and laboratory characteristics of the multisystem inflammatory syndrome in children: a case series of 75 patients. *Arch Pediatr Infect Dis.* 2022;10(4):e120863.
- Jamalidoust M, Ashkan Z, Pouladfar G, et al. Prevalence and clinical presentation of COVID 19 in health care workers in two main hospitals during the pandemic in Shiraz, Iran. *Arch Pediatr Infect Dis.* 2022;10(4):e121753.
- Pourakbari B, Mahmoudi S, Mahmoudieh Y, et al. SARS-CoV-2 RNAemia in children: an Iranian referral hospital-based study. *J Med Virol.* 2021;93(9):5452-5457.
- Mertz C, Glowinski R, Cohen SH, et al. Severe acute respiratory syndrome coronavirus 2 RNAemia and clinical outcomes in children with coronavirus disease 2019. *J Infect Dis.* 2022;225(2):208-213.
- Armin S, Mirkarimi M, Pourmoghaddas Z. Evidence-based prediction of COVID-19 severity in hospitalized children. *Int J Clin Pract.* 2022;2022:1918177.
- Sena GR, Lima T, Vidal SA, et al. Clinical characteristics and mortality profile of COVID-19 patients aged less than 20 years old in pernambuco - Brazil. *Am J Trop Med Hyg.* 2021;104(4):1507-1512.
- Kang S-J, S IJL. Jung, and chemotherapy age-related morbidity and mortality among patients with COVID-19. *Infect Chemother.* 2020;52(2):154-164.