

RESEARCH

Open Access



The effect of distance nurse-led fatigue management on fatigue, sleep quality, and self-efficacy in patients with multiple sclerosis: a quasi-experimental study

Mahla Qomi¹, Mahnaz Rakhshan² , Mohsen Ebrahimi Monfared³ and Zahra Khademian^{4*}

Abstract

Background Fatigue is one of the most common problems in patients with multiple sclerosis (MS) and has adverse effects on their sleep status and self-efficacy. This study aimed to determine the effect of distance nurse-led fatigue management on fatigue, sleep quality, and self-efficacy in patients with MS.

Methods This quasi-experimental study was performed on 60 patients with MS in Arak, Iran. Subjects were randomly assigned into intervention and control groups. The intervention group received eight sessions of nurse-led fatigue management training through the Skyroom platform. The control group received only the usual programs. Data were collected before and two months after the intervention using the Fatigue Severity Scale, the Pittsburgh Sleep Quality Index, and the Multiple Sclerosis Self-Efficacy Scale. The significance level in this study was determined 0.05.

Results After the intervention, the mean score of fatigue severity in the intervention group was significantly lower than the control group (2.52 ± 0.40 vs 5.65 ± 0.52) ($P < 0.001$). Also, after the intervention, the mean score of self-efficacy in the intervention group was significantly higher than the control group (49.37 ± 3.25 vs 24.43 ± 2.52) ($P < 0.001$). Furthermore, after the intervention the mean score of sleep quality was lower in intervention group (11.92 ± 2.01) than the control group (15.46 ± 1.40) ($P < 0.001$).

Conclusion Distance nurse-led fatigue management improved fatigue, sleep quality, and self-efficacy in patients with MS. We recommend the use of these courses as an important step toward improving fatigue, sleep quality, and self-efficacy among these patients.

Keywords Fatigue, Multiple sclerosis, Nursing care, Patient care management, Self efficacy, Sleep quality, Telenursing

Background

Multiple sclerosis (MS) is the most common progressive neurologic disease in young adults worldwide [1]. According to a report in 2020, about 2.8 million people worldwide are living with this disease [2]. Iran is a country in the Middle East with moderate to high prevalence of MS, ranging from 5.30 to 74.28 per 100,000 individuals [3, 4]. Fatigue is one of the most common symptoms of MS, ranged from 36.5 to 78.0%. It can significantly affect the patients' quality of life and impose an economic

*Correspondence:

Zahra Khademian
zahrakhademian@yahoo.com; khademian@sums.ac.ir

¹ Student Research Committee, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran

² Community Based Psychiatric Care Research Center, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran

³ Department of Neurology, School of Medicine, Arak University of Medical Sciences, Arak, Iran

⁴ Department of Nursing, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Conclusion

The findings of this study showed that the distance nurse-led fatigue management can improve fatigue, sleep quality, and self-efficacy in patients with MS. These findings showed the important role of nurses in community-oriented care of patients with MS. Since nurses can play an effective role in the management of patients with MS, they should pay more attention to the patients' fatigue. Moreover, it is necessary for them to learn fatigue reduction skills and include them in patient care protocols. We recommend teaching these skills to nursing students and including these skills in the nursing continuing education. In addition, we recommend the inclusion of nurse-led fatigue management as an inexpensive non-pharmacological intervention in the management of patients with MS. The distance nature of similar interventions provides the possibility of access to the people in remote areas, as well as providing services to patients in situations such as the COVID-19 pandemic. We recommend further research with a longer follow-up period on the effectiveness of similar interventions.

Abbreviations

ANCOVA	Analysis of Covariance
COVID-19	Coronavirus Disease of 2019
MS	Multiple Sclerosis
MSSES	Multiple Sclerosis Self-Efficacy Scale
PSQI	Pittsburgh Sleep Quality Index
FSS	Fatigue Severity Scale
SPSS	Statistical Package for the Social Sciences

Acknowledgements

This manuscript was extracted from the Master's thesis by Mahla Qomi, which was sponsored by the Vice Chancellor for Research of Shiraz University of Medical Sciences, Shiraz, Iran. Hereby, appreciation goes to the Vice Chancellor for Research of Shiraz University of Medical Sciences for its financial support. The authors would like to thank Clinical Research Development Center of Nemazee Hospital, Dr. Nasrin Shokrpour for editorial assistance, and Mr Ali Mohammad Keshavarz and Dr Parvin Ghaem Maghami for their assistance in data analysis. We would also like to thank the respectable authorities and personnel of the MS Association, Arak, and all patients who earnestly helped us with this project.

Authors' contributions

MQ, ZKh, MR, and MEM made substantial contributions to the conception and design of the study. Data was collected by MQ. Data analysis and interpretation were done by ZKh, MQ, and MR. MQ conducted the intervention. ZKh and MQ participated in drafting the manuscript. ZKh, MQ, MR, and MEM revised the manuscript critically for important intellectual content and final approval of the manuscript.

Funding

The present study was financially supported by the Vice Chancellor for Research of Shiraz University of Medical Sciences, Shiraz, Iran (Grant No. 22080). The funding body did not play any roles in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

Data resource and statistical analysis outputs can be provided by the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The approval of the study was obtained from the Ethics Committee of Shiraz University of Medical Sciences (ethics code: IR.SUMS.REC.1399.1035) and the officials of the MS Association. Furthermore, all methods were carried out in accordance with Declaration of Helsinki and relevant guidelines and regulations. The goals and process of the research were also explained to the patients. All participants signed the informed consent form. Participation in the study was voluntary and participants could withdraw at any time. In addition, the participants' information was kept confidential.

Consent to publication

Not applicable.

Competing interest

The authors declare no competing interests.

Received: 18 August 2022 Accepted: 9 February 2023

Published online: 14 February 2023

References

1. Deuschl G, Beghi E, Fazekas F, Varga T, Christoforidi KA, Sipido E, Bassetti CL, Vos T, Feigin VL. The burden of neurological diseases in Europe: an analysis for the Global Burden of Disease Study 2017. *Lancet Public Health*. 2020;5:e551–67.
2. Walton C, King R, Rechtman L, Kaye W, Leray E, Marrie RA. Rising prevalence of multiple sclerosis worldwide: insights from the Atlas of MS, third edition. *Mult Scler*. 2020;26:1816–21.
3. Sahraian MA, Sahebkar S, Dehghani R, Derakhshan Jazari M, Kazami Moghaddam V, Kouchaki E. Multiple sclerosis-a disease on a dramatically rising trend in Iran: Review of possible reasons. *Curr J Neurol*. 2017;16(1):34–40.
4. Asmarian N, Sharafi Z, Mousavi A, Jacques R, Tamayo I, Bind M-A, Abutorabi-zarchi M, Moradian MJ, Izadi S. Multiple sclerosis incidence rate in southern Iran: a Bayesian epidemiological study. *BMC Neurol*. 2021;21:309. <https://doi.org/10.1186/s12883-021-02342-1>.
5. Oliva Ramirez A, Keenan A, Kalau O, Worthington E, Cohen L, Singh S. Prevalence and burden of multiple sclerosis-related fatigue: a systematic literature review. *BMC Neurol*. 2021;21:1–16.
6. Harrison AM, Safari R, Mercer T, Picariello F, van der Linden ML, White C, Moss-Morris R, Norton S. Which exercise and behavioural interventions show most promise for treating fatigue in multiple sclerosis? a network meta-analysis. *Mult Scler*. 2021;27:1657–78.
7. Zielińska-Nowak E, Włodarczyk L, Kostka J, Miller E. New strategies for rehabilitation and pharmacological treatment of fatigue syndrome in multiple sclerosis. *J Clin Med*. 2020;9:3592.
8. Phyto AZZ, Demaneuf T, De Livera AM, Jelinek GA, Brown CR, Marck CH, Neate SL, Taylor KL, Mills T, O'Kearney E, Karahalios A, Weiland TJ. The efficacy of psychological interventions for managing fatigue in people with multiple sclerosis: a systematic review and meta-analysis. *Front Neurol*. 2018;9:149. <https://doi.org/10.3389/fneur.2018.00149>.
9. Kotterba S, Neusser T, Norenberg C, Bussfeld P, Glaser T, Dörner M, Schürks M. Sleep quality, daytime sleepiness, fatigue, and quality of life in patients with multiple sclerosis treated with interferon beta-1b: results from a prospective observational cohort study. *BMC Neurol*. 2018;18:123.
10. Veauthier C, Gaede G, Radbruch H, Wernecke KD, Paul F. Sleep disorders reduce health-related quality of life in multiple sclerosis (Nottingham health profile data in patients with multiple sclerosis). *Int J Mol Sci*. 2015;16:16514–28.
11. Barun B. Pathophysiological background and clinical characteristics of sleep disorders in multiple sclerosis. *Clin Neurol Neurosurg*. 2013;115(Suppl 1):S82–85.
12. Aljundi NA, Kelly M, Zeineddine S, Salloum A, Pandya N, Shamim-Uzzaman QA, Badr AN, Mitchell MN, Sankari A, Badr MS, Martin JL. Sleep disorders, daytime symptoms and quality of life in veterans with multiple sclerosis: preliminary findings. *Sleep Adv*. 2022;3(1):012.