

Determinant of antibacterial failure in patients with head and neck infections of odontogenic origin

Hamed Gheibollahi¹, Sona Mousavi², Hossein Daneste², Mohammad Mehdi Taheri², Ali Peyravi³, Amirhossein Babaei^{4,5}

¹Department of Oral and Maxillofacial Surgery, School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

²Department of Oral and Maxillofacial Surgery, School of Dentistry, Shahid Rajaei Acute Care Surgical Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

³Student Research Committee, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran

⁴Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran

⁵Otolaryngology Research Center, Department of Otolaryngology, Shiraz University of Medical Sciences, Shiraz, Iran

Article history: Received: 29.09.2022 Accepted: 31.01.2023 Published: 03.02.2023

ABSTRACT:

Introduction: Odontogenic infections are polymicrobial in origin and can be life-threatening. Antibacterial failure is an important issue in the treatment of odontogenic infections.

Aim: This study aimed to determine the factors associated with antibacterial failure in patients with head and neck infections of odontogenic origin.

Material and method: This retrospective epidemiological study was performed using data collected from 229 patients with head and neck infections of odontogenic origin who were operated on in Shahid Rajaei hospital from March 2014 to December 2019.

Results: 123 (53.7%) patients were female and there were 106 (46.3%) males. The mean age $\pm$ SD was 33.01 ± 13.37 years (range 7 to 80). The most common offending teeth were lower molars (81.7%) and lower premolars (5.7%). The most common site of infection was the submandibular area (36.4%) followed by the buccal (20.4%) and pterygomandibular (17.1%) regions. The most common pathogen was *Streptococcus haemolyticus*. The length of hospitalization was higher (4.66 days) in patients with failure of treatment compared to those without it (6.00 days) ($P = 0.002$). A combination of penicillin G and metronidazole was prescribed for all patients with failure of treatment compared to 57.6% in patients without failure of treatment ($P = 0.002$). There was no statistically significant difference between the two groups regarding age, duration of illness before hospitalization, WBC, gender, history of chemotherapy, hypertension, smoking, pregnancy, alcohol consumption, diabetes mellitus, the rate of fever, trismus, dysphagia, malaise, antibiotic before hospitalization, or surgical approach.

Conclusion: Possible determinants in this study were not associated with antibacterial failure. Further studies should be conducted to investigate this relationship.

KEYWORDS:

antibacterial failure, Iran, odontogenic infections

ABBREVIATIONS

CT – computed tomography

SD – standard deviation

WBC – white blood cell

INTRODUCTION

Odontogenic infections affect the tooth and its supporting structures and are polymicrobial in origin; they are caused by aerobic and anaerobic microorganisms [1]. Despite huge advances in medical sciences, many patients still suffer from odontogenic infections in the world [2]. Rapid documentation of infections and appropriate treatment are critical due to life-threatening complications like airway obstruction, sepsis, necrotizing fasciitis,

cavernous sinus thrombosis, and mediastinitis [3]. The most prevalent presentation of patients admitted due to odontogenic head and neck infections was trismus, toxic appearance, and dysphagia [4].

The source of odontogenic infection could be dental caries, gingival and periodontal disease. Many odontogenic infections are self-limiting. Though, some of them may spread into the adjacent spaces and cause more serious infections [5]. Antimicrobial resistance is an increasing global healthcare concern and rapid diagnosis of microorganisms is crucial [6]. Broad-spectrum antibiotics are widely used to treat infections [7]. There is no consensus between different countries on treatment protocols, and intravenous penicillin G, metronidazole, clindamycin, amoxicillin-clavulanic acid, or amoxicillin-sulbactam are the most commonly prescribed antibiotics [8].

The present study has some strengths. We gathered data from a large group of patients from a referral centre. This increases the external validity of data. On the other hand, there are some limitations such as the lack of follow-up data as it was a retrospective study. We suggest further studies with a prospective design.

CONCLUSION

Based on information obtained in our analysis, possible determinants

in this study were not associated with antibacterial failure. Further prospective studies should be conducted to investigate this relationship.

ACKNOWLEDGEMENT

The authors would like to thank Shiraz University of Medical Sciences, Shiraz, Iran and also the Center for Development of Clinical Research of Nemazee Hospital and Dr Nasrin Shokrpour for editorial assistance.

REFERENCES

- López-González E., Vitales-Noyola M., González-Amaro A.M. et al.: Aerobic and anaerobic microorganisms and antibiotic sensitivity of odontogenic maxillofacial infections. *Odontology*, 2019; 107(3): 409–417.
- Haque M., Sartelli M., Haque S.Z.: Dental Infection and Resistance-Global Health Consequences. *Dent J (Basel)*, 2019; 7(1): 22.
- Bali R.K., Sharma P., Gaba S., Kaur A., Ghanghas P.: A review of complications of odontogenic infections. *Natl J Maxillofac Surg.*, 2015; 6(2): 136–143.
- Aliabadi E., Farshad M.M., Kheirkhah M., Jafari S.H.: Clinical, CT scan, and laboratory changes of abscess patients with odontogenic origin admitted to Shiraz acute surgical care center, Iran. *Fam. Med. Prim. Care Rev.*, 2021; 10(9): 3314–3318.
- Caruso S.R., Yamaguchi E., Portnof J.E.: Update on Antimicrobial Therapy in Management of Acute Odontogenic Infection in Oral and Maxillofacial Surgery. *Oral Maxillofac Surg Clin North Am.*, 2022; 34(1): 169–177.
- Babaei A.H., Pouladfar G., Pourabbas B. et al.: Seven-Year Trend of Antimicrobial Resistance of *Acinetobacter* and *Pseudomonas* spp. Causing Bloodstream Infections: A Retrospective Study from Shiraz, Southern Iran. *Jundishapur J. Microbiol.*, 2019; 12(4): e85819.
- Babaei H., Motamedifar M., Khalifat S. et al.: In vitro study of antibacterial property and cytotoxic effects of aqueous, ethanolic, methanolic, and hydroalcoholic extracts of fenugreek seed. *Pak. J. Med.*, 2018; 12(2): 906–910.
- Jevon P., Abdelrahman A., Pigadas N.: Management of odontogenic infections and sepsis: an update. *Br Dent J.*, 2020; 229(6): 363–370.
- Yuvaraj V.: Maxillofacial Infections of Odontogenic Origin: Epidemiological, Microbiological and Therapeutic Factors in an Indian Population. *Indian J. Otolaryngol.*, 2016; 68(4): 396–399.
- Eshghpour M., Sabouri M., Labafchi A., Shooshtari Z., Samieirad S.: Maxillofacial Infections: A 5-Year Assessment of the Epidemiology and Etiology in an Iranian Population. *J. Oral Maxillofac. Surg.*, 2021; <https://doi.org/10.1007/s12663-021-01569-2>.
- Katoumas K., Anterriotis D., Fyrgiola M. et al.: Epidemiological analysis of management of severe odontogenic infections before referral to the emergency department. *J Craniomaxillofac Surg.*, 2019; 47(8): 1292–1299.
- Uittamo J., Löfgren M., Hirvikangas R., Furuholm J., Snäll J.: Severe odontogenic infections: focus on more effective early treatment. *Br J Oral Maxillofac Surg.*, 2020; 58(6): 675–680.
- Neal T.W., Hammad Y., Carr B.R., Schlieve T.: The Cost of Surgically Treated Severe Odontogenic Infections: A Retrospective Study Using Severity Scores. *J Oral Maxillofac Surg.*, 2022; 80(5): 897–901. doi: 10.1016/j.joms.2022.01.001.
- Keswani E.S., Venkateshwar G.: Odontogenic Maxillofacial Space Infections: A 5-Year Retrospective Review in Navi Mumbai. *J. Oral Maxillofac. Surg.*, 2019; 18(3): 345–353.
- Shukla A., Mehrotra D.: Odontogenic Infections: General Principles. *Oral Maxillofac Surg.*, 2021; 429–439. doi: https://doi.org/10.1007/978-981-15-1346-6_20.
- Weise H., Naros A., Weise C., Reinert S., Hoefert S.: Severe odontogenic infections with septic progress – a constant and increasing challenge: a retrospective analysis. *BMC Oral Health.*, 2019; 19(1): 173.
- Bahl R., Sandhu S., Singh K., Sahai N., Gupta M.: Odontogenic infections: Microbiology and management. *Contemp. Clin. Dent.*, 2014; 5(3): 307–311.
- Lim S.W., Lee W.S., Mani S.A., Kadir K.: Management of odontogenic infection in paediatric patients: a retrospective clinical study. *Eur Arch Paediatr Dent.*, 2020; 21(1): 145–154.
- Qu L., Liang X., Jiang B. et al.: Risk Factors Affecting the Prognosis of Descending Necrotizing Mediastinitis From Odontogenic Infection. *J. Oral Maxillofac. Surg.*, 2018; 76(6): 1207–1215.
- Ogle O.E.: Odontogenic Infections. *Dent. Clin. N. Am.*, 2017; 61(2): 235–252.
- Rastenienė R., Pūrienė A., Aleksejūnienė J., Pečiulienė V., Zaleckas L.: Odontogenic Maxillofacial Infections: A Ten-Year Retrospective Analysis. *Surg Infect.*, 2015; 16(3): 305–312.
- Umeshappa H., Shetty A., Kavatagi K. et al.: Microbiological profile of aerobic and anaerobic bacteria and its clinical significance in antibiotic sensitivity of odontogenic space infection: A prospective study of 5 years. *Natl J Maxillofac Surg.*, 2021; 12(3): 372–379.