

RESEARCH

Open Access



The possible relationship between the healthy eating index-2015 and the 10-year risk of cardiovascular diseases

Pegah Ahmadijoo¹, Mohammad Hassan Eftekhari^{1*}, Seyed Jalil Masoumi^{2,3}, Maryam Ranjbar Zahedani⁴ and Farzaneh Mohammadi^{1,5}

Abstract

Background Cardiovascular diseases (CVDs) remain the leading cause of mortality worldwide. This underlies the need to evaluate different targets, such as diet quality. In this regard, we conducted the present study to find whether the healthy eating index-2015 (HEI-2015) score is associated with a 10-year risk of CVDs based on Framingham Risk Score (FRS) and QRISK3 in different body mass index (BMI) groups.

Methods This cross-sectional study was performed based on Shiraz University of Medical Sciences Employees Health Cohort Study (SUMS EHCS) data in April 2020. A total of 764 participants met the inclusion criteria. An expert performed demographic, anthropometric, and dietary evaluations. A semi-quantitative food frequency questionnaire (FFQ) was applied to assess the diet quality, and FRS and QRISK3 were used to evaluate the 10-year risk of CVDs.

Results Based on the results, many components of HEI-2015 indicated an increasing trend through quartiles ($p < 0.001$). However, the consumption of refined grains in higher quartiles showed a decreasing trend ($p < 0.001$). The consumption of added sugar and saturated fatty acids (SFAs) in higher quartiles revealed an increasing trend ($p < 0.001$). In addition, lower HEI-2015 scores and lower whole grain consumption were significantly associated with higher BMI ($p < 0.05$). Also, lower consumption of fruits showed a significant relationship with higher risk scores of Framingham and QRISK3 ($p < 0.05$). Higher added sugar and SFAs intake was significantly related to lower FRS ($p < 0.05$). A significant reverse association between HEI-2015 and QRISK3 and Framingham risk scores was seen ($p < 0.05$).

Conclusion Our findings support dietary recommendations to increase fruit and whole grains intake to prevent CVD and obesity. Moreover, a significant inverse association between HEI-2015 and QRISK3 and Framingham risk scores was observed. Since the results for added sugars and SFA intakes were controversial, further studies are needed.

Keywords Body mass index, Cardiovascular disease, Framingham risk score, QRISK3, Healthy eating index

*Correspondence:

Mohammad Hassan Eftekhari

h_eftekhari@yahoo.com; eftekhari@sums.ac.ir

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Abbreviations

CVDs	Cardiovascular diseases
WHO	World health organization
FRS	Framingham risk score
HEI	Healthy eating index
SUMS EHCS	Shiraz university of medical sciences employees health cohort study
BMI	Body mass index
HDL	High-density lipoprotein cholesterol
LDL	Low-density lipoprotein cholesterol
FFQ	Food frequency questionnaire
SFA	Saturated fatty acid
PUFA	Polyunsaturated fatty acid
MUFA	Monounsaturated fatty acid
SD	Standard deviation
IQR	Interquartile range
CHD	Coronary heart disease

Acknowledgements

The authors would like to thank Shiraz University of Medical Sciences, Shiraz, Iran, and also the Center for Development of Clinical Research of Nemazee Hospital and Dr. Nasrin Shokrpour for editorial assistance.

Authors' contributions

Mohammad Hassan Eftekhari conceived and supervised the study. Pegah Ahmadijoo analyzed and interpreted the data. Seyed Jalil Masoumi conducted and supervised the study and data collecting. Pegah Ahmadijoo, Maryam Ranjbar Zahedani, and Farzaneh Mohammadi wrote the manuscript. All the authors critically revised the manuscript.

Funding

This study was financially supported by the Vice-Chancellor of Research and Technology of Shiraz University of Medical Sciences, Shiraz, Iran. The funding body played no role in the design of the study and collection, analysis, interpretation of data, and in writing the manuscript.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of Shiraz University of Medical Sciences, Shiraz, Iran (Code: IR.SUMS.REC.1399.482). The study was also conducted in accordance with the Helsinki Declarations of Ethics. The informed consent form was completed for all the patients.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran. ²Nutrition Research Center, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran. ³Gastroenterohepatology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. ⁴Department of Nutrition Sciences, School of Health, Larestan University of Medical Sciences, Larestan, Iran. ⁵Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran.

Received: 21 January 2023 Accepted: 22 June 2023

Published online: 28 June 2023

References

- Wang H, Naghavi M, Allen C, Barber RM, Bhutta ZA, Carter A, Casey DC, Charlson FJ, Chen AZ, Coates MM. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388:1459–544.
- Balakumar P, Maung-U K, Jagadeesh G. Prevalence and prevention of cardiovascular disease and diabetes mellitus. *Pharmacol Res*. 2016;113:600–9.
- Sarrafadegan N, Mohammadifard N. Cardiovascular disease in Iran in the last 40 years: prevalence, mortality, morbidity, challenges and strategies for cardiovascular prevention. *Arch Iran Med*. 2019;22:204–10.
- Damen JA, Pajouheshnia R, Heus P, Moons KG, Reitsma JB, Scholten RJ, Hoof L, Debray TP. Performance of the Framingham risk models and pooled cohort equations for predicting 10-year risk of cardiovascular disease: a systematic review and meta-analysis. *BMC Med*. 2019;17:1–16.
- Dhaliwal SS, Welborn TA. Central obesity and multivariable cardiovascular risk as assessed by the Framingham prediction scores. *Am J Cardiol*. 2009;103:1403–7.
- Elagizi A, Kachur S, Carbone S, Lavie CJ, Blair SN: A Review of Obesity, Physical Activity, and Cardiovascular Disease. *Curr Obes Rep*. 2020;9:1–11.
- Im E, Kim G-S. Relationship between sleep duration and Framingham cardiovascular risk score and prevalence of cardiovascular disease in Koreans. *Medicine*. 2017;96:e7744.
- Ghosal S, Sinha B, Ved J, Biswas M. Quantitative measure of asymptomatic cardiovascular disease risk in Type 2 diabetes: evidence from Indian outpatient setting. *Indian Heart J*. 2020;72:119–22.
- Aa A. Diet quality concept. *Nutrition*. 2014;30:613–8.
- Nicklas TA, O'Neil CE, Fulgoni VL III. Diet quality is inversely related to cardiovascular risk factors in adults. *J Nutr*. 2012;142:2112–8.
- Tande DL, Magel R, Strand BN. Healthy Eating Index and abdominal obesity. *Public Health Nutr*. 2010;13:208–14.
- Gesteiro E, Bernal BR, Bastida S, Sánchez-Muniz F. Maternal diets with low healthy eating index or Mediterranean diet adherence scores are associated with high cord-blood insulin levels and insulin resistance markers at birth. *Eur J Clin Nutr*. 2012;66:1008–15.
- Xu Z, Steffen LM, Selvin E, Rebholz CM. Diet quality, change in diet quality, and risk of incident cardiovascular disease and diabetes. *Public Health Nutr*. 2020;23:329.
- Esmailinezhad Z, Babajafari S, Sohrabi Z, Eskandari M-H, Amooee S, Barati-Boldaji R. Effect of synbiotic pomegranate juice on glycemic, sex hormone profile and anthropometric indices in PCOS: a randomized, triple blind, controlled trial. *Nutr Metab Cardiovasc Dis*. 2019;29:201–8.
- Preston SH, Fishman E, Stokes A. Effects of categorization and self-report bias on estimates of the association between obesity and mortality. *Ann Epidemiol*. 2015;25(907–911):e902.
- Khanna NN, Jamthikar AD, Gupta D, Nicolaidis A, Araki T, Saba L, Cuadrado-Godia E, Sharma A, Omerzu T, Suri HS. Performance evaluation of 10-year ultrasound image-based stroke/cardiovascular (CV) risk calculator by comparing against ten conventional CV risk calculators: a diabetic study. *Comput Biol Med*. 2019;105:125–43.
- Ghayour-Mobarhan M, Moohebbati M, Esmaily H, Ebrahimi M, Parizadeh SMR, Heidari-Bakavoli AR, Safarian M, Mokhber N, Nematy M, Saber H. Mashhad stroke and heart atherosclerotic disorder (MASHAD) study: design, baseline characteristics and 10-year cardiovascular risk estimation. *Int J Public Health*. 2015;60:561–72.
- Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol*. 1985;122:51–65.
- Rahmani J, Varkaneh HK, Ryan PM, Zarezaadeh M, Rashvand S, Clark C, Day AS, Hekmatdoost A. Healthy Eating Index-2015 as a predictor of ulcerative colitis risk in a case-control cohort. *J Dig Dis*. 2019;20:649–55.
- Esmaily Z, Tajary Z, Daei S, Rezaei M, Eyzvakhani A, Motlagh ARD, Palmowski A. Association between Healthy Eating Index-2015 scores and probable sarcopenia in community-dwelling Iranian older adults: a cross-sectional study. *J Nutr Sci*. 2021;10:e20.