



THE EFFECT OF ORAL PROTEIN SUPPLEMENTATION ON THE GROWTH OF VERY LOW BIRTH WEIGHT PRETERM INFANTS ADMITTED TO THE NEONATAL INTENSIVE CARE UNIT: A RANDOMIZED CLINICAL TRIAL

Fariba Hemmati¹ , Maral Ghassemzadeh^{2*} 

¹Associate Professor of Pediatrics and Neonatologist, Neonatal Research Center, Department of pediatrics, School of Medicine, Shiraz University of Medical Science, Shiraz, Iran

²Assistant Professor of Neonatology, Tehran University of Medical Sciences, Hakim Children Hospital, Tehran, Iran

Abstract

Background: During NICU admission, extra-uterine growth retardation that can affect the neurodevelopmental outcome is a challenging problem in extremely preterm infants. This trial aimed to determine the effect of additional enteral protein supplementation on the growth velocity of the anthropometric parameters.

Method: In this randomized controlled trial, 77 preterm infants (gestational age ≤ 33 weeks and birth weight < 1500 g) who reached full enteral feeding with either fortified breast milk or preterm formula were included. They were randomized to receive either 4- < 5 g/kg/day protein through extra protein supplementation (intervention) or 3- < 4 g/kg/day protein. Weight gain, as well as length and head circumference growth, were monitored daily and weekly, respectively. Venous blood gas, blood urea nitrogen (BUN), and albumin levels were checked weekly.

Results: Five out of 77 participants were excluded due to feeding intolerance. Analyses were conducted on 36 neonates with protein intake of 3.66 ± 0.22 gr/kg/day and 36 with extra protein intake. Baseline characteristics were similar between the groups. An additional protein supply of 0.89 gr/kg/day, resulting in an average protein intake of 4.55 ± 0.18 in the intervention group, increased the postnatal weight gain, linear growth, and head circumference growth (7.98 gr/kg/day, 0.347 cm/week, and 0.38 cm/week, respectively). The albumin levels were significantly increased, but the BUN levels were not significantly increased in the intervention group. None of the patients developed necrotizing enterocolitis or significant acidosis.

Conclusion: Protein supplementation significantly improves the growth of the anthropometric parameters. An increase in serum albumin and no increase in serum urea can indicate the anabolic effect of extra protein. Protein supplementation can add to routine feeding protocols of VLBW infants without any short-term adverse effect; however, further study for evaluation of long-term complications is needed.

Keywords

Protein supplement, Preterm infants, Growth, Nutrition, VLBW

Received: 2022 November 20

Accepted: 2023 January 12

Introduction

The last trimester of pregnancy, which is often lost in preterm birth, is the time when the fetus has the most protein intake for growth. Furthermore, extremely preterm infants lose protein approximately twofold that of term infants [1]. Therefore,

preterm newborns often experience a negative nitrogen balance. Prematurity is a nutritional emergency since studies demonstrate that these infants experience extra-uterine growth retardation as a result of a cumulative deficiency in

*Corresponding author: Maral Ghassemzadeh, M.D.
e-mail: maral.ghassemzadeh1985@gmail.com

Copyright © 2023 Hemmati and Ghassemzadeh.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 Public License (CC BY) (<https://creativecommons.org/licenses/by/4.0/>). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

albumin levels which were significantly higher in the protein supplement group and increased significantly in this group throughout the study (based on repeated measures analysis). In the Cochrane review published at 2020 [24] which evaluated high versus low protein intake in the formula-fed LBW infants, no study addressed serum albumin as a secondary outcome; however, in studies which compared very high versus low protein intake, Cooke et al. [25] reported serum albumin levels and found no significant differences between the two groups. Van der Aker et al. reported that albumin synthesis was stimulated by parenterally administered amino acid in preterm infants [26]. The increase in serum albumin in our neonates can indicate the anabolic effect of extra protein and increased weight gain velocity is due to protein and fat accretion. One putative risk for high protein intake is increased concentrations of hydrogen ions as a result of the immaturity of amino acid metabolic pathways in preterm infants [27], and metabolic acidosis may develop. Hence, we evaluated the acid-base status of the participants. The PH was significantly lower, while PCO₂ and HCO₃ were higher in the protein group than the controls (after intervention). Based on repeated measures analysis, however, the changes in the PH level and HCO₃ levels were not statistically significant throughout the study in both groups. Thus, no significant metabolic acidosis developed, like the result of Cooke et al.'s study [25].

Our study had some limitations such as a lack of long-term follow-up of our patients and the small sample size. However, based on group matchings and insignificant differences between the baseline features of our groups, we believe that our study could provide insight into further multicenter RCTs and review of articles to obtain the most satisfactory treatment and supplement regimen for neonates.

Conclusion

Protein supplement significantly improved the growth of all aspects of the anthropometric parameters including weight, length, and head circumference. No short-term adverse effect was detected during the study. This study could provide insight into further multicenter RCTs to define the optimal levels of protein intake for VLBW neonates and add protein supplement to the routine feeding protocols. Furthermore, long-term studies are needed to evaluate long-term benefits, especially on neurodevelopment and probable complications of this intervention.

Conflicts of interest

No conflicts of interest were declared by the authors.

Acknowledgments

The present article was extracted from the thesis written by Maral Ghassemzadeh and was financially supported by Shiraz University of Medical Sciences with the grant number of 95-01-01-13884. The authors would like to thank the Office of Vice-chancellor for Research at Shiraz University of Medical Sciences for the financial support. The authors would also like to thank Shiraz University of Medical Sciences, Shiraz, Iran and also Center for Development of Clinical Research of Nemazi Hospital and Dr. Nasrin Shokrpour for editorial assistance.

Abbreviations

GA: Gestational Age
 BUN: Blood Urea Nitrogen
 VLBW: Very Low Birth Weight
 NEC: Necrotizing Enterocolitis
 SGA: Small for Gestational Age
 HC: Head Circumference
 FBM: Fortified Breast Milk
 PF: Premature Formula
 PNGF: Post Natal Growth Failure
 NICDH: National Institute of Child Health and Human Development
 PDA: Patent Ductus Arteriosus
 MOM: Mothers' Own Milk
 AAP: American Academy of Pediatrics
 ESPGHAN: European Society for Pediatric Gastroenterology Hepatology and Nutrition

ORCID ID's:

Fariba Hemmati 0000-0002-5704-3947 (<https://orcid.org/0000-0002-5704-3947>)

Maral Ghassemzadeh 0000-0002-3407-8119 (<https://orcid.org/0000-0002-3407-8119>)

Author Contributorship (CRediT Roles):

Fariba Hemmati: Conceptualization and supervision and main idea. Maral Ghassemzadeh: Data collection, analysis, writing the manuscript.

Funding

The present article was extracted from the thesis written by Maral Ghassemzadeh and was financially supported by Shiraz University of Medical Sciences with the grant number of 95-01-01-13884.