

RESEARCH

Open Access



Margin status impact on recurrence of phyllodes tumors in high-risk groups: a retrospective observational study

Aliyeh Ranjbar¹, Vahid Zangouri^{1,2*} and Mansoureh Shokripour^{3*}

Abstract

Background Phyllodes tumor (PT) is an fibroepithelial tumor with potential for local recurrence. The optimal margin for surgical resection of PT is still debated, particularly in cases of positive margins. This study aimed to identify the risk factors for phyllodes tumor recurrence and the effect of a free margin on tumor recurrence by considering these risk factors.

Materials and methods This is a retrospective observational study of patients diagnosed with PT who had undergone surgical management. The data were collected from medical records from 2001 to 2020 in the breast clinic of Shahid Motahhari Clinic of Shiraz. Patients were followed up for at least 3 years after the operation to be checked for local recurrence or distant metastasis at regular intervals.

Results This retrospective study included 319 patients with PT who underwent surgical management. Of these patients, 83.9% ($n = 267$), 7.6% ($n = 24$), and 8.5% ($n = 27$) were classified as benign, borderline, and malignant, respectively. 8.8% of all patients and 7.6% of non-malignant cases experienced local recurrence, and risk factors for recurrence included oral contraceptive use, smoking, size > 4 cm, stromal overgrowth, and stromal cell atypia. A negative surgical margin decreased the prevalence of recurrence in tumors > 4 cm and with stromal overgrowth significantly.

Conclusion The study found that a negative margin in all patients did not reduce the recurrence rate in benign and borderline phyllodes tumors, suggesting close follow up as a reasonable alternative. However, a negative margin may be effective in reducing recurrence in certain high-risk groups.

Keywords Phyllodes tumour, Local recurrence, Surgical margins, Risk factor

*Correspondence:

Vahid Zangouri
zangouri.vahid1390@gmail.com
Mansoureh Shokripour
Shokripurm@sums.ac.ir

¹Breast Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

²Surgical Oncology Division, General Surgery Department, Shiraz University of Medical Sciences, Shiraz, Iran

³Department of Pathology, Medical School, Shiraz University of Medical Sciences, Shiraz, Iran



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 3 Kaplan-Meier survival plot of recurrence-free survival. stratified by the negative and positive surgical margin in mass size > 4 cm (A), and present of stromal overgrowth (B)

Acknowledgements

This study was extracted from the dissertation which was written by Aliyeh Ranjbar for the degree of Doctor of Medicine (thesis no. 20039). The authors would like to thank Shiraz University of Medical Sciences, Shiraz, Iran and also Center for Development of Clinical Research of Nemazee Hospital and Dr. Nasrin Shokrpour for editorial assistance.

Author contributions

V.Z., A.R., and M.Sh.: conceptualized the data. M.Sh., and A.R.: curated the data. A.R.: wrote the original draft. V.Z.: edited the draft. All authors reviewed the manuscript.

Funding

The project was funded by Vice Chancellor for Research of the Shiraz University of Medical Science [Grant No. 99-01-01-22758].

Data availability

Data is available from the corresponding author upon reasonable request via email.

Declarations

Ethics approval and consent to participate

This study was conducted in accordance with the principles established by the Declaration of Helsinki and obtained the approval of the Ethics Committee of Shiraz University of Medical Sciences (approval ID: IRSUMS.MED.REC.1400.029). Informed consent was obtained and signed by all study participants prior to recruitment.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 14 August 2023 / Accepted: 28 December 2023

Published online: 09 January 2024

References

- Reinfuss M, Mituś J, Duda K, Stelmach A, Ryś J, Smolak K. The treatment and prognosis of patients with phyllodes tumor of the breast: an analysis of 170 cases. *Cancer: Interdisciplinary International Journal of the American Cancer Society*. 1996;77(5):910–6.
- Guerrero MA, Ballard BR, Grau AM. Malignant phyllodes tumor of the breast: review of the literature and case report of stromal overgrowth. *Surg Oncol*. 2003;12(1):27–37.
- Guillot E, Couturaud B, Reyal F, Curnier A, Ravinet J, Laé M, et al. Management of phyllodes breast tumors. *Breast J*. 2011;17(2):129–37.
- Bennett IC, Khan A, De Freitas R, Chaudary MA, Millis RR. Phyllodes tumours: a clinicopathological review of 30 cases. *Aust N Z J Surg*. 1992;62(8):628–33. <https://doi.org/10.1111/j.1445-2197.1992.tb07534.x>. Epub 1992/08/01. PubMed PMID: 1322660.
- The world Health Organization Histological Typing of Breast Tumors–Second Edition. The World Organization. *Am J Clin Pathol*. 1982;78(6):806–16. <https://doi.org/10.1093/ajcp/78.6.806>. PubMed PMID: 7148748.
- Zhang Y, Kleer CG. Phyllodes Tumor of the Breast: Histopathologic Features, Differential Diagnosis, and Molecular/Genetic Updates. *Arch Pathol Lab Med*. 2016;140(7):665–71. Epub 2016/07/01. <https://doi.org/10.5858/arpa.2016-0042-RA>. PubMed PMID: 27362571.
- Yilmaz E, Sal S, Lebe B. Differentiation of phyllodes tumors versus fibroadenomas: mammographic and sonographic features. *Acta Radiol*. 2002;43(1):34–9.
- Michaud P, Chave B, Lemaire B, Maitre F, Tescher M. Les tumeurs phyllodes du sein. *Rev Fr Gynecol Obstet*. 1989;84(12):944–9.
- Liberman L, Bonaccio E, Hamele-Bena D, Abramson AF, Cohen MA, Dershaw DD. Benign and malignant phyllodes tumors: mammographic and sonographic findings. *Radiology*. 1996;198(1):121–4.
- Vos D, Mastboom W, De Vos R. Phyllodes tumor in the breast. *Ned Tijdschr Geneesk*. 1998;142(13):716–21.
- Iau P, Lim T, Png D, Tan W. Phyllodes tumour: an update of 40 cases. *Ann Acad Med Singapore*. 1998;27(2):200–3.
- Shet T, Rege J. Cystic degeneration in phyllodes tumor. A source of error in cytologic interpretation. *Acta Cytol*. 2000;44(2):163–8.
- Mangi AA, Smith BL, Gadd MA, Tanabe KK, Ott MJ, Souba WW. Surgical management of phyllodes tumors. *Arch Surg*. 1999;134(5):487–92. <https://doi.org/10.1001/archsurg.134.5.487>. PubMed PMID: 10323420.
- Rowell MD, Perry RR, Hsui JG, Barranco SC. Phyllodes tumors. *Am J Surg*. 1993;165(3):376–9. [https://doi.org/10.1016/s0002-9610\(05\)80849-9](https://doi.org/10.1016/s0002-9610(05)80849-9). PubMed PMID: 8383473.
- Sallusti E, Simonelli I, Marzullo A, Peis A, Modesti M. [The breast phyllodes tumor: surgical therapy following histological transformation. Case Report] *G Chir*. 2004;25(3):95–7. Epub 2004/07/01. PubMed PMID: 15224663.
- Gabriele R, Borghese M, Corigliano N, Barbaro M, Conte M. Phyllodes tumor of the breast. Personal contribution of 21 cases. *Il Giornale Di Chirurgia*. 2000;21(11–12):453–6.
- Mishra SP, Tiwary SK, Mishra M, Khanna AK. Phyllodes tumor of breast: a review article. *ISRN Surg*. 2013;2013:361469. <https://doi.org/10.1155/2013/361469>. PubMed PMID: 23577269; PubMed Central PMCID: PMC3615633.
- Mangi AA, Smith BL, Gadd MA, Tanabe KK, Ott MJ, Souba WW. Surgical management of phyllodes tumors. *Arch Surg*. 1999;134(5):487–93.
- Kılıç MÖ, Terzioğlu SG, Bozkurt B, Dağlar G. Phyllodes Tumor of the breast: analysis of 48 patients. *J Breast Health*. 2016;12(4):158–64. <https://doi.org/10.5152/tjbh.2016.3100>. PubMed PMID: 28331755.
- Moo T-A, Alabdulkareem H, Tam A, Fontanet C, Lu Y, Landers A, et al. Association between recurrence and re-excision for close and positive margins versus observation in patients with benign phyllodes tumors. *Ann Surg Oncol*. 2017;24(10):3088–92.
- Ouyang Q, Li S, Tan C, Zeng Y, Zhu L, Song E, et al. Benign phyllodes tumor of the breast diagnosed after ultrasound-guided vacuum-assisted biopsy: surgical excision or wait-and-watch? *Ann Surg Oncol*. 2016;23:1129–34.
- Rosenberger LH, Thomas SM, Nimbkar SN, Hieken TJ, Ludwig KK, Jacobs LK, et al. Contemporary multi-institutional cohort of 550 cases of phyllodes tumors (2007–2017) demonstrates a need for more individualized margin guidelines. *J Clin Oncol*. 2021;39(3):178.