



Journal of Nutrition and Food Security

Shahid Sadoughi University of Medical Sciences
School of Public Health
Department of Nutrition



Shahid Sadoughi
University of Medical Sciences
School of Public Health
Nutrition Department

eISSN: 2476-7425

pISSN: 2476-7417

JNFS 2024; 9(2): 251-264

Website: jnfs.ssu.ac.ir

Comparing of Metabolic Syndrome Components, Inflammation, Cortisol Level, and Psychological Distress in Obese/Overweight and Normal Weight Women

Golazin Hoseini; MSc¹, Esmail Soltani; PhD², Najmeh Hejazi; PhD^{*3}, Maedeh Gordali; MSc⁴ & Zahra Sohrabi; PhD⁵

¹ Student Research Committee, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran; ² Psychiatry and Behavioral Sciences Research Center, Shiraz University of Medical Sciences, Shiraz, Iran; ³ Nutrition Research Center, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran; ⁴ Student Research Committee, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran; ⁵ Nutrition Research Center, Department of Nutrition Community, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran.

ARTICLE INFO

ORIGINAL ARTICLE

Article history:

Received: 10 May 2022

Revised: 26 Sep 2022

Accepted: 26 Sep 2022

*Corresponding author

najmehhejazi@gmail.com
Department of Clinical
Nutrition, School of
Nutrition and Food sciences,
Shiraz University of Medical
Sciences, Shiraz, Iran.

Postal code: 7153675500

Tel: +98 917 7020859

ABSTRACT

Background: High levels of stress in obese people, hypothalamic-pituitary-adrenal (HPA) axis disorder, and social pressures can increase cortisol level and lead to psychological disorders. The aim of this study is to compare psychological distress, biochemical parameters, and metabolic syndrome components between normal-weight and overweight (OW)/obese women. **Methods:** This was an analytical cross-sectional study conducted on 75 women aged 18 to 60; they were divided into three groups: obese and OW on diet (obese/OW on diet), obese and OW without diet (obese/OW without diet), and normal-weight. The components of metabolic syndrome, serum cortisol, and high sensitive C-reactive protein (hs-CRP) levels were measured. General health questionnaire-28 (GHQ-28) was also completed to assess psychological distress. **Result:** The results revealed that there was a significant difference between normal weight and the other two groups regarding metabolic syndrome components, which included waist circumference (WC), fasting blood sugar (FBS), systolic blood-pressure (SBP), and hs-CRP ($P < 0.05$). Serum cortisol level was significantly higher in obese/OW on diet compared with the other two groups ($P < 0.001$). Moreover, the total-GHQ score was significantly lower in normal weight compared with the group of obese/OW on diet and the group without diet ($P = 0.001$). **Conclusion:** Being on a diet may expose a person to stress and increase the serum cortisol level. Elevated psychological distress, metabolic syndrome components, and inflammation were apparent in obese and OW women compared to normal-weight ones.

Keywords: Obesity; Metabolic syndrome; Inflammation; Psychological distress; Diet; Non-communicable diseases

Introduction

Obesity, as a chronic multifactorial disease, results from a positive balance of energy and excess fat accumulation throughout life. Obesity leads to structural abnormalities, physiological

damage, and functional disorders when left untreated (Jastreboff *et al.*, 2019). According to the World Health Organization (WHO), overweight and obesity are among the top five causes of

This paper should be cited as: Hoseini G, Soltani E, Hejazi N, Gordali M, Sohrabi Z. Comparing of Metabolic Syndrome Components, Inflammation, Cortisol Level, and Psychological Distress in Obese/Overweight and Normal Weight Women. *Journal of Nutrition and Food Security (JNFS)*, 2024; 9(2): 251-264.

Conclusion

Going on a diet, irrespective of weight changes, might expose a person to stress and increase the serum cortisol levels; a significant weight loss was probably required to improve the components of metabolic syndrome and inflammation. It was also found that psychological distress, metabolic syndrome components and inflammation were higher in obese and OW cases compared with normal-weight ones.

Acknowledgements

This study was financially supported by the Vice Chancellor of Research and Technology at Shiraz University of Medical Sciences, Shiraz, Iran (grant no: 97-01-84-19277). The authors would like to thank the participants, for their cooperation, Shiraz University of Medical Sciences, the Center for Development of Clinical Research of Nemazee Hospital for their support, and Dr. Nasrin Shokrpour for his editorial assistance.

Conflict of interests

The authors declared no conflict of interest.

Authors' Contribution

Soltani E, Hejazi N, Sohrabi Z and Gordali M wrote and reviewed the manuscript; Soltani E was involved with the research method, ; Hejazi N was the managed the project, and Gordali M analyzed data. All the authors approved the final manuscript.

Funding

This study was financially supported by the Vice Chancellor of Research and Technology Department at Shiraz University of Medical Sciences, Shiraz, Iran (grant no: 97-01-84-19277).

References

- Aadahl M & Jørgensen T 2003. Validation of a new self-report instrument for measuring physical activity. *Medicine and science in sports and exercise*. **35** (7): 1196-1202.
- Abraham S, Rubino D, Sinaii N, Ramsey S & Nieman L 2013. Cortisol, obesity, and the metabolic syndrome: A cross-sectional study of obese subjects and review of the literature. *Obesity*. **21** (1): E105-E117.
- Akter R, et al. 2017. Effect of Obesity on Fasting Blood Sugar. *Mymensingh medical journal*. **26** (1): 7-11.
- Al-Safi ZA, et al. 2018. Evidence for disruption of normal circadian cortisol rhythm in women with obesity. *Gynecological endocrinology*. **34** (4): 336-340.
- Alimoradi Z, et al. 2020. Weight-related stigma and psychological distress: A systematic review and meta-analysis. *Clinical nutrition*. **39** (7): 2001-2013.
- Amatruda JM, Livingston JN & Lockwood DH 1985. Cellular mechanisms in selected states of insulin resistance: human obesity, glucocorticoid excess, and chronic renal failure. *Diabetes/metabolism reviews*. **1** (3): 293-317.
- Atlantis E & Ball K 2008. Association between weight perception and psychological distress. *International journal of obesity*. **32** (4): 715-721.
- Barazzoni R, Gortan Cappellari G, Ragni M & Nisoli E 2018. Insulin resistance in obesity: an overview of fundamental alterations. *Eating and weight disorders-studies on anorexia, bulimia and obesity*. **23** (2): 149-157.
- Björntorp P 1996. The regulation of adipose tissue distribution in humans. *Journal of the international association for the study of obesity*. **20** (4): 291-302.
- Björntorp P & Rosmond R 2000. Obesity and cortisol. *Nutrition*. **16** (10): 924-936.
- Black S, Kushner I & Samols D 2004. C-reactive protein. *Journal of biological chemistry*. **279** (47): 48487-48490.
- Bujang MA, Sa'at N & Bakar TMITA 2017. Determination of minimum sample size requirement for multiple linear regression and analysis of covariance based on experimental and non-experimental studies. *Epidemiology, biostatistics, and public health*. **14** (3): e12117-12111.
- Burhans MS, Hagman DK, Kuzma JN, Schmidt KA & Kratz M 2018. Contribution of adipose tissue inflammation to the development of type 2 diabetes mellitus. *Comprehensive physiology*. **9** (1): 1.
- Chan J, Sauve B, Tokmakejian S, Koren G & Van Uum S 2014. Measurement of cortisol and