

CASE REPORT

Open Access



A middle-aged man with dyspnea and hoarseness: an unusual case of vocal cord paralysis

Seyedeh Nadia Tabatabaeifar¹, Mahsa Ahmadpour² and Mohammad Javad Fallahi^{3*}

Abstract

Background Pleuroparenchymal fibroelastosis (PPFE) is a rare and distinct form of Interstitial lung disease predominantly affecting the upper lung zones. It is characterized by fibrotic thickening of the visceral pleura and adjacent subpleural parenchyma. While common complications include spontaneous pneumothorax and pneumomediastinum, vocal cord paralysis (VCP) or paresis has been increasingly recognized as a potential manifestation in recent reports.

Case presentation We present a 49-year-old man presenting with progressive dyspnea, hoarseness, and left-sided vocal cord paralysis. Imaging studies revealed upper lobe-dominant fibrotic changes associated with significant pleural thickening consistent with PPFE. A comprehensive evaluation ruled out secondary causes of PPFE and other potential etiologies of VCP. Despite supportive management, the patient's VCP persisted, likely due to architectural distortion of the lung affecting the recurrent laryngeal nerve pathway.

Conclusions This case adds to the limited but growing body of literature on the association between PPFE and VCP. Understanding this rare complication is crucial for early recognition and appropriate management, as it highlights the diverse clinical manifestations of PPFE and its impact on patient outcomes.

Keywords Pleuroparenchymal fibroelastosis, Vocal cord paralysis, Interstitial lung disease, Recurrent laryngeal nerve, Tracheobronchial distortion

Background

Pleuroparenchymal (PPFE) is a rare form of interstitial pneumonia characterized by an upper lung zone-predominant fibrotic process involving the visceral pleura and adjacent subpleural parenchyma [1]. While the disease is largely idiopathic, non-idiopathic cases of PPFE have increasingly been reported in association with other interstitial lung diseases, including idiopathic pulmonary fibrosis, hypersensitivity pneumonitis (HP), and familial pulmonary fibrosis [1]. In clinical practice, although the definitive diagnosis relies on pathological evidence, this may not always be feasible due to the disease's rapid progression and the risk of respiratory complications

*Correspondence:

Mohammad Javad Fallahi
falahijavad@yahoo.com

¹Internal Medicine Department, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

²Shahid Faghihi Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

³Thoracic and Vascular Surgery Research Center, Shiraz University of Medical Sciences, Shiraz, Iran



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



Fig. 2 Left vocal cord paralysis without mucosal abnormalities on fiberoptic bronchoscopy

Table 1 Summary of patient's body plethysmography and DLCO. FEV1 (liter): forced expiratory volume in one second, FVC (liter): forced vital capacity, TLC (liter): total lung capacity, RV (liter): residual volume, DLCO (mmol/kpa/minute): lung diffusion capacity for carbon monoxide

Parameter	Predicted	Actual	Percent %
FEV1	4.00	1.51	38
FVC	5.16	1.70	33
FEV1/FVC		0.84	
TLC	7.46	4.89	66
RV	2.17	3.10	143
RV/TLC	32	63	199
DLCO	11.16	5.86	52

Discussion

The primary etiologies of VCP involve conditions that affect the recurrent laryngeal nerve (RLN), including tumors of the thyroid, esophagus, mediastinum, and pleura; tuberculosis; iatrogenic causes after cervical or mediastinal surgery; diabetes mellitus; mitral valve stenosis; and rapid swelling of the pulmonary artery. In rare instances, some paranchymal lung disease such as coal workers' pneumoconiosis with massive fibrosis, lung collapse due to cystic fibrosis, and invasive pulmonary aspergillosis, idiopathic or secondary mediastinal fibrosis may also result in VCP [4–9]. Typically, the left RLN is more often affected, as its path is longer, as observed in this case [10, 11].

Although the mechanism of RLN paralysis in lung diseases remains unclear, recent reports propose several mechanisms. These include stretching or retraction of the RLN due to chest wall adhesion resulting from pleural fibrosis [7], distortion of the tracheobronchial tree leading to RLN traction or compression [3, 11], and left-sided RLN compression at the aortopulmonary window, where the aortic arch and left pulmonary artery are

in close proximity [3, 7, 11]. Lardinois et al. reported a case of left RLN paralysis associated with silicosis, where progressive recovery of the voice was observed 15 weeks after careful dissection of the nerve and release from scar encasement during video mediastinoscopy [6].

To date, seven cases of PPFE with VCP have been reported, including the present case. In all but one case, the paralysis involved the left vocal cord [11]. Dominant left RNP along right tracheal deviation, support RLN traction or compression as most probable mechanism of vocal cord paralysis in PPFE. After excluding treatable and more dangerous cause of new onset hoarseness, VCP secondary to parenchymal involvement should be considered in PPFE patient.

Clinicians encountering patients with upper lobe-predominant fibrotic lung disease and concurrent VCP should include PPFE as a differential diagnosis.

Abbreviations

PPFE	Pleuroparenchymal fibroelastosis
VCP	Vocal cord paralysis
IPPFE	Idiopathic pleuroparenchymal fibroelastosis
RLN	Recurrent laryngeal nerve
HP	Hypersensitivity pneumonitis
IIPs	Idiopathic interstitial pneumonias
CT	Computed tomography
MRI	Magnetic resonance imaging
VLS	Video laryngeal stroboscopy
PCR	Polymerase chain reaction
RV	Residual volume
TLC	Total lung capacity
BAL	Bronchoalveolar lavage

Acknowledgements

The authors would like to thank Shiraz University of Medical Sciences, Shiraz, Iran and also Center for Development of Clinical Research of Nemazee Hospital and Dr. Nasrin Shokrpour for editorial assistance.

Author contributions

SNT analyzed and interpreted the patient's clinical data and contributed to drafting the manuscript. MA assisted with patient management, performed the literature review, and contributed to manuscript preparation. MJF