


Protective effects of limb remote ischemic per-conditioning on the heart injury induced by renal ischemic-reperfusion through the interaction of the apelin with the RAS/iNOS pathway

Sahar Janfeshan, Fatemeh Masjedi, Zeinab Karimi*

Shiraz Nephro-Urology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Article Info



Article Type:
Original Article

Article History:

Received: 14 Aug. 2022
 Revised: 28 Jan. 2023
 Accepted: 13 Mar. 2023
 ePublished: 8 Oct. 2023

Keywords:

Limb remote ischemic per-conditioning,
 Renal ischemia-reperfusion,
 Myocardial injury,
 Apelin,
 Renin-angiotensin system,
 iNOS


Abstract

Introduction: Remote ischemic conditioning upregulates endogenous protective pathways in response to ischemia-reperfusion injury. This study tested the hypothesis that limb remote ischemic per-conditioning (RIPerC) exerts cardioprotective effects via the renin-angiotensin system (RAS)/inducible nitric oxide synthase (iNOS)/apelin pathway.

Methods: Renal ischemia-reperfusion injury (I/R) was induced by bilateral occlusion of the renal pedicles for 60 minutes, followed by 24 hours of reperfusion; sham-operated rats served as controls. RIPerC was induced by four cycles (5 minutes) of limb ischemia-reperfusion along with bilateral renal ischemia. The functional disturbance was evaluated by renal (BUN and creatinine) and cardiac (troponin I and lactate dehydrogenase) injury biomarkers.

Results: Renal I/R injury increased renal and cardiac injury biomarkers that were reduced in the RIPerC group. Histopathological findings of the kidney and heart were also suggestive of amelioration injury-induced changes in the RIPerC group. Assessment of cardiac electrophysiology revealed that RIPerC ameliorated the decline in P wave duration without significantly affecting other cardiac electrophysiological changes. Further, renal I/R injury increased the plasma (322.40 ± 34.01 IU/L), renal (8.27 ± 1.10 mIU/mg of Protein), and cardiac (68.28 ± 10.28 mIU/mg of protein) angiotensin-converting enzyme (ACE) activities in association with elevations in the plasma and urine nitrite (25.47 ± 2.01 & 16.62 ± 3.05 $\mu\text{mol/L}$) and nitrate (15.47 ± 1.33 & 5.01 ± 0.96 $\mu\text{mol/L}$) levels; these changes were reversed by RIPerC. Further, renal ischemia-reperfusion injury significantly ($P=0.047$) decreased the renal (but not cardiac) apelin mRNA expression, while renal and cardiac ACE2 ($P<0.05$) and iNOS ($P=0.043$) mRNA expressions were significantly increased compared to the sham group; these effects were largely reversed by RIPerC.

Conclusion: Our results indicated that RIPerC protects the heart against renal ischemia-reperfusion injury, likely via interaction of the apelin with the RAS/iNOS pathway.



Introduction

Renal ischemia-reperfusion (I/R) is a major cause of acute kidney injury (AKI) that is usually induced in some surgeries (kidney transplantation and coronary artery bypass grafting) and multiple clinical situations, including trauma, severe infection, sepsis, and hemorrhagic

shock.^{1,2} Renal I/R is characterized by sudden restriction of blood supply to the kidney followed by the subsequent restoration of perfusion and re-oxygenation.^{3,4}

The consequences of renal I/R injury are local as well as remote organ destruction such as liver,⁵ lung,^{6,7} and heart.⁸ The physiological and pathophysiological



*Corresponding author: Zeinab Karimi, Email: zkarimi@sums.ac.ir



© 2024 The Author(s). This work is published by BioImpacts as an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

Research Highlights

What is the current knowledge?

✓ Remote ischemic conditioning upregulates endogenous protective pathways in response to ischemia-reperfusion injury.

✓ The physiological and pathophysiological interaction between the kidney and the cardiovascular system provides essential indicators for maintaining homeostasis.

What is new here?

✓ Limb remote ischemic per conditioning improves the myocardium dysfunction induced by renal ischemia-reperfusion injury through the interaction of the apelin with the RAS/iNOS pathway.

diseases.⁵⁵ Apelin depletion progressed the myocardium dysfunction and structural remodeling through angiotensin II pathways.⁵⁶ Our data clearly indicated that remote ischemic per-conditioning suppressed the cardiac ACE activity and ACE2 expression.

Conclusion

Remote ischemic per-conditioning is a potential phenomenon to promote endogenous protective pathways against renal I/R injury. The findings of the current study revealed that renal I/R induced kidney and heart structural and functional disturbance by a significant increase in injury biomarkers. Histological damages in the renal and heart tissues were in line with the increased ACE activity and NO metabolites in the tissues and biological samples. On the other hand, the level of apelin mRNA was downregulated, but iNOS and ACE2 genes were upregulated in the renal and heart tissues. In total, our results indicated that RIPC protects the heart against renal I/R injury, probably through the interaction of the apelin with the RAS/iNOS pathway.

Acknowledgments

The authors would like to thank the Center for Development of Clinical Research of Nemazee Hospital, Shiraz, Iran, and Dr. Nasrin Shokrpour for editorial assistance.

Authors' Contribution

Conceptualization: Zeinab Karimi.

Data curation: Sahar Janfeshan, Fatemeh Masjedi.

Formal analysis: Fatemeh Masjedi.

Funding acquisition: Zeinab Karimi.

Investigation: Fatemeh Masjedi, Zeinab Karimi.

Methodology: Sahar Janfeshan, Fatemeh Masjedi.

Project administration: Zeinab Karimi.

Resources: Zeinab Karimi.

Software: Fatemeh Masjedi.

Supervision: Zeinab Karimi.

Validation: Zeinab Karimi.

Visualization: Zeinab Karimi.

Writing—original draft: Sahar Janfeshan, Fatemeh Masjedi.

Writing—review & editing: Zeinab Karimi.

Competing interests

The authors declare no conflict of interest.

Ethical Statement

The experimental protocols were approved by the local ethics committee of Shiraz University of Medical Sciences (Approval ID: IR.SUMS.REC.1399.894).

Data Availability Statement

Data reported in this manuscript are available upon reasonable request from the corresponding author.

Funding

This study was supported by the Vice-Chancellor for Research, Shiraz University of Medical Sciences (Academic Grant Number: 21395).

References

- Ronco C, Bellomo R, Kellum JA. Acute kidney injury. *Lancet* **2019**; 394: 1949-64. [https://doi.org/10.1016/S0140-6736\(19\)32563-2](https://doi.org/10.1016/S0140-6736(19)32563-2)
- Roozbeh J, Malekmakan L, Monavarian M, Daneshian A, Karimi Z. Survival of Kidney Retransplant Compared With First Kidney Transplant: A Report From Southern Iran. *Exp Clin Transplant* **2018**; 16: 386-90. <https://doi.org/10.6002/ect.2016.0130>
- Malek M, Nematbakhsh M. Renal ischemia/reperfusion injury; from pathophysiology to treatment. *J Renal Inj Prev* **2015**; 4: 20-7. <https://doi.org/10.12861/jrip.2015.06>
- Karimi Z, SoukhakLari R, Rahimi-Jaberi K, Esmaili Z, Moosavi M. Nanomicellar curcuminoids attenuates renal ischemia/reperfusion injury in rat through prevention of apoptosis and downregulation of MAPKs pathways. *Mol Biol Rep* **2021**; 48: 1735-43. <https://doi.org/10.1007/s11033-021-06214-2>
- Serteser M, Koken T, Kahraman A, Yilmaz K, Akbulut G, Dilek ON. Changes in Hepatic TNF- α Levels, Antioxidant Status, and Oxidation Products after Renal Ischemia/Reperfusion Injury in Mice. *J Surg Res* **2002**; 107: 234-40. <https://doi.org/10.1006/jsre.2002.6513>
- Karimi Z, Ketabchi F, Alebrahimdehkordi N, Fatemikia H, Owji SM, Moosavi SM. Renal ischemia/reperfusion against nephrectomy for induction of acute lung injury in rats. *Ren Fail* **2016**; 38: 1503-15. <https://doi.org/10.1080/0886022x.2016.1214149>
- Hashemi SS, Janfeshan S, Karimi Z. Acute lung injury induced by acute uremia and renal ischemic-reperfusion injury: The role of toll-like receptors 2 and 4, and oxidative stress. *Iran J Basic Med Sci* **2022**; 25: 643-51. <https://doi.org/10.22038/ijbms.2022.64025.14099>
- El-Sayed SS, Shahin RM, Fahmy A, Elshazly SM. Quercetin ameliorated remote myocardial injury induced by renal ischemia/reperfusion in rats: Role of Rho-kinase and hydrogen sulfide. *Life Sci* **2021**; 287: 120144. <https://doi.org/10.1016/j.lfs.2021.120144>
- Amann K, Wanner C, Ritz E. Cross-Talk between the Kidney and the Cardiovascular System. *J Am Soc Nephrol* **2006**; 17: 2112-9. <https://doi.org/10.1681/ASN.2006030204>
- Kelly KJ. Distant Effects of Experimental Renal Ischemia/Reperfusion Injury. *J Am Soc Nephrol* **2003**; 14: 1549-58. <https://doi.org/10.1097/01.ASN.0000064946.94590.46>
- Nath KA, Norby SM. Reactive oxygen species and acute renal failure. *Am J Med* **2000**; 109: 665-78. [https://doi.org/10.1016/S0002-9343\(00\)00612-4](https://doi.org/10.1016/S0002-9343(00)00612-4)
- Fatemikia H, Ketabchi F, Karimi Z, Moosavi SM. Distant effects of unilateral renal ischemia/reperfusion on contralateral kidney but not lung in rats: the roles of ROS and iNOS. *Can J Physiol Pharmacol* **2016**; 94: 477-87. <https://doi.org/10.1139/cjpp-2015-0285>
- Efrati S, Berman S, Hamad RA, Siman-Tov Y, Ilgiyaev E, Maslyakov I, et al. Effect of captopril treatment on recuperation from ischemia/reperfusion-induced acute renal injury. *Nephrol Dial Transplant* **2012**; 27: 136-45. <https://doi.org/10.1093/ndt/gfr256>
- Panico K, Abrahão MV, Trentin-Sonoda M, Muzi-Filho H, Vieyra