







The effect of damask rose aromatherapy on anxiety and pain in endoscopic lithotripsy patients: A Double-blind Randomized Clinical Trial

Zinat Mohebbi¹ , Maryam Mohit^{2*} , Fatemeh Shirazi¹ , Parvin Ghaemmaghami³ 

1. Nursing Department, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran

2. Surgical Technology Department, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran

3. School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran

* Correspondence: Maryam Mohit, Surgical Technology Department, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran.

Tel: +989305046290; Email: MaryamMohit71@gmail.com

Abstract

Background: Aromatherapy is a modality within the realm of alternative medicine to manage pain and alleviate anxiety in high-stress circumstances. The primary objective of this study was to examine the impact of damask rose on the levels of anxiety and pain experienced by patients under endoscopic lithotripsy.

Methods: The present double-blind randomized clinical trial was conducted at hospitals in Shiraz, Iran, in 2022, involving 120 patients undergoing endoscopic lithotripsy. These patients were randomly allocated into control and intervention groups using a permuted block design. The data collection was done using the Spielberger and the Visual Analog Scale questionnaires at three stages: the baseline period, the pre-operative holding area, and six hours following the surgical procedure. The intervention group was exposed to a cotton swab infused with three drops of damask rose for a duration of 30 minutes. The data analysis included descriptive statistics, repeated measure analysis of variance, and pairwise comparisons utilizing the Bonferroni post hoc test.

Results: The mean levels of overt anxiety in the intervention group during the first and second phases of the intervention were 46.17 ± 7.18 ($P < 0.001$) and 46.32 ± 3.24 ($P = 0.021$). Furthermore, the mean covert anxiety levels in the intervention group throughout the first and second phases were found to be 45.10 ± 7.83 ($P = 0.003$) and 45.87 ± 3.59 ($P < 0.001$). The intervention group exhibited a mean pain level of 5.7 ± 1.01 during the initial phase and 2.53 ± 0.81 throughout the second phase ($P < 0.001$).

Conclusion: The findings of this study indicated that rose aromatherapy reduced both overt and covert anxiety levels, as well as the pain experienced by patients both pre- and post-endoscopic lithotripsy.

Article History

Received: 14 January 2024

Received in revised form: 5 May 2024

Accepted: 19 June 2024

Published online: 26 June 2024

DOI: [10.29252/jgbrnm.21.2.11](https://doi.org/10.29252/jgbrnm.21.2.11)

Keywords

Anxiety

Pain

Aromatherapy

Lithotripsy

Endoscopy

Article Type: Original Article



© The author(s)

Highlights

What is current knowledge?

According to the existing literature, no study has assessed effect of damask rose aromatherapy on the patients' anxiety and pain before and after endoscopic lithotripsy.

What is new here?

This study shows that the use of inhalation damask rose aromatherapy results in temporary pain alleviation and a decrease in both overt and covert anxiety levels among individuals following endoscopic lithotripsy.

Introduction

Kidney stones represent a prevalent urological condition, necessitating treatment through a range of procedures, including transurethral lithotripsy (TUL), percutaneous nephrolithotomy (PCNL), and extracorporeal shock wave lithotripsy (ESWL). Presently, the aforementioned treatment modality has emerged as the primary therapeutic option for managing sizable kidney stones in the renal pelvic, Calix, and calyceal diverticulum (1).

In contemporary surgical practice, there has been a notable shift towards employing minimally invasive techniques in the treatment of patients across all medical disciplines. This trend is particularly pronounced in the realm of urological treatments pertaining to urinary system stones (2). The endoscopic approach has been shown to effectively decrease postoperative pain. This strategy has several advantages, including the reduction of hospitalization duration, cost reduction, and enhanced patient satisfaction. While endoscopic surgery has been shown to mitigate postoperative pain by minimizing tissue manipulation and utilizing smaller incisions, it is important to acknowledge that no surgical procedure is entirely devoid of pain and anxiety. Consequently, several approaches are employed to alleviate the pain and anxiety (3).

Surgery is commonly considered a stressful experience for both the patients and their family (4). Surgical anxiety has been found to induce alterations in the physiological functioning of the human body, hence posing potential risks to an individual's overall health and impeding the process of tissue recovery and repair (5). Anxiety has been seen to result in several physiological and behavioral manifestations. These include diminished appetite, discomfort, sleep

disturbances, compromised immune function, delayed wound healing, elevated vital signs, impaired concentration, decreased engagement in self-care tasks, as well as lower collaboration with healthcare professionals (6).

The presence of anxiety has been found to impact the level of postoperative pain experienced and the subsequent healing process following anesthesia (7). Effective management of anxiety during the preoperative phase has the potential to promote cardiovascular stability, thus enhancing the patient's overall health throughout the surgical procedure and subsequent recovery period. Additionally, the management of anxiety has been found to be associated with a decrease in the use of anesthetics during surgical procedures, improved pain tolerance, and faster patient discharge from medical facilities (8).

Postoperative pain is among the most common and important issues of patient care. A significant majority of patients, over 80%, encounter immediate postoperative pain following surgical procedures, with about 70% of patients reporting pain levels ranging from moderate to severe (9,10). Recently, medication therapy is no longer regarded as the optimal and exclusive approach for mitigating the patients' pain and anxiety. While pharmaceuticals effectively alleviate the patient's pain and anxiety, they frequently cause undesirable adverse reactions (11). In this context, the utilization of many approaches in supplementary medicine might prove advantageous. Complementary and alternative medicine (CAM) therapies encompass a range of strategies employed for disease prevention, treatment, and the enhancement of overall health and well-being in conjunction with conventional medical interventions. A widely embraced kind of alternative medicine that has gained significant popularity is aromatherapy, which involves the utilization of various aromatic substances (12, 13).

Damask rose is a medicinal plant whose essential oil is utilized in the practice of aromatherapy (14). It has been reported that rose has sedative, analgesic, antioxidant, antibacterial, and antidiabetic effects (15). Iran is recognized as one of the leading nations in rose production globally, which is well-known as Mohammadi flower, placing it in the top four countries in this industry (16). The purpose of this study is to investigate the potential benefits of aromatherapy, specifically the inhalation of Damask rose, on anxiety and pain experienced by patients undergoing endoscopic lithotripsy. Despite the increasing popularity of aromatherapy, there remains a limited number of studies that have examined its effectiveness in this context, highlighting the need for further investigation. Our research team aims to address this gap in the literature and provide valuable insights into the potential role of aromatherapy in improving patient outcomes in this population.

Discussion

The present study was conducted to examine the impact of rose aromatherapy on the levels of anxiety and pain experienced by patients undergoing endoscopic lithotripsy. The findings of this study indicated that within the intervention group, the damask rose aromatherapy resulted in a significant reduction in the mean pain throughout the initial and second phases of the intervention, as compared to the baseline period. Furthermore, the results of the present study indicated that the intervention group experienced a reduction in the mean levels of overt and covert anxieties during both stages of aromatherapy, as compared to the control group.

Sadeghi et al. (2020) conducted a study to investigate the impact of damask rose aromatherapy on the level of pain experienced by burn patients. The findings of their study revealed a significant reduction in pain intensity among patients following the intervention (18). A meta-analysis conducted by Nasiri et al. (2021) examined the impact of damask rose aromatherapy on the severity of acute pain in adults. The study revealed a significant decrease in pain intensity among patients following the intervention (22). The study conducted by Bastani and colleagues in 2017 demonstrated that using damask rose essence in aromatherapy can have a positive impact on pain relief following knee arthroscopic surgery (23). Additionally, a study by Roozbahani and team in 2015 found that inhalation of damask rose water can help reduce labor pain ($p < 0.001$), which supports the findings of the current study (24).

A clinical trial by Abbasi et al. (2020) examined the impact of damask rose aromatherapy on the severity of postoperative pain, the overt and covert anxiety levels experienced by women after a caesarean section. The researchers employed damask rose essential oil as an intervention for a duration of 30 minutes among the participants. The researchers demonstrated a significant reduction in maternal pain levels, and also the mean score of overt and covert anxiety following surgery was significantly lower in the intervention group compared to the control group (4), which supports the findings of the current study. Moreover, in line with the present study, Bikmoradi et al. examined the impact of damask rose aromatherapy on anxiety levels among patients undergoing coronary angiography. The study demonstrated a statistically significant difference in the mean anxiety score in the intervention group compared with the control group (25).

Farzaneh et al. (2022) investigated the impact of damask rose aromatherapy on the levels of overt and covert anxiety among patients scheduled for lithotripsy operations prior to the procedure. A statistically significant difference ($P < 0.001$) was seen when they compared the mean levels of overt and covert anxiety between the control and intervention groups. They revealed a reduction in overt and covert anxiety levels among patients in the intervention group prior to surgery (26). Also, Mirzaee et al. conducted a study that demonstrated the effectiveness of Damask rose essential oil aromatherapy in reducing pre-endoscopic anxiety among patients (27).

Fazlollahpour et al.'s study aimed to examine the impact of inhaling rose essential oil through aromatherapy on anxiety levels in patients undergoing coronary artery bypass graft. The findings of this study indicated that, in contrast to the current study, there was no significant reduction in the severity of anxiety observed among the intervention groups (28). This lack of significant effect may potentially be attributed to the duration of exposure to the rose aroma. Azizi et al. (2018) conducted a study that demonstrated a reduction in pain before and after dressing through the inhalation of lavender essence. Furthermore, there was no meaningful variance in pain intensity between the placebo and control groups. The primary distinction between the current study and the earlier one is the inhaler type used (29). The research conducted in this study was focused on patients who underwent lithotripsy procedures. The study has several limitations, including the reluctance of some patients to cooperate in the research, the absence of a dedicated space for implementing aromatherapy, and the lack of comparable studies on damask rose inhalation in this patient population. Consequently, it was not feasible to draw comparisons with existing research findings. As a result, prospective studies should focus on replicating this research with a different demographic and patients with varied medical conditions, exploring diverse aromas, and investigating the optimal duration of aromatherapy.

Conclusion

The use of inhalation damask rose aromatherapy was found to result in temporary pain alleviation and a decrease in both overt and covert anxiety levels among individuals following endoscopic lithotripsy. Given the cost-effectiveness of this intervention for patients, nurses can employ it as a kind of individualized treatment to attain relaxation and secure its short-term advantages.

Acknowledgement

This project resulted from an MSc thesis in Surgical Technology Major at Shiraz University of Medical Sciences. The authors would like to thank Shiraz University of Medical Sciences, Shiraz, Iran and also the Center for Development of Clinical Research of Nemazee Hospital and Dr Nasrin Shokrpour for editorial assistance.

Funding sources

The study was supported by Shiraz University of Medical Sciences (Grant No. 26218).

Ethical statement

The current study was approved by the research ethics committee of Shiraz University of Medical Sciences (IR.SUMS.NUMIMG.REC.1401.064). Additionally, it has been registered in the Iranian Registry of Clinical Trials (IRCT) with the registration code of IRCT20100919004775N14.

Conflicts of interest

The authors declare that they have no competing interests.

Author contributions

All authors participated in the conception and design of the study. MM and ZM contributed to proposal preparation and data collection. MM and PGH participated in data analysis and interpretation. MM, FSH and ZM contributed to drafting of the article. All authors have seen and approved the submitted version of this manuscript.

References

- Baralo B, Samson P, Hoenig D, Smith A. Percutaneous kidney stone surgery and radiation exposure: A review. *Asian J Urol.* 2020;7(1):10-7. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Hernandez N, Mozafarpour S, Song Y, Eisner BH. Cessation of ureteral colic does not necessarily mean that a ureteral stone has been expelled. *J Urol.* 2018;199(4):1011-4. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Noh GO, Park KS. Effects of aroma self-foot reflexology on peripheral neuropathy, peripheral skin temperature, anxiety, and depression in gynaecologic cancer patients undergoing chemotherapy: A randomised controlled trial. *Eur J Oncol Nurs.* 2019;42:82-9. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Abbasijahromi A, Hojati H, Nikooei S, Jahromi HK, Dowlatkhan HR, Zarean V, et al. Compare the effect of aromatherapy using lavender and Damask rose essential oils on the level of anxiety and severity of pain following C-section: A double-blinded randomized clinical trial. *J Complement Integr Med.* 2020;17(3). [View at Publisher] [DOI] [PMID] [Google Scholar]
- Rahmati H, Seidi J, Ghodsbin F, Rahimi S, Gholamvaysi B. The effect of music therapy on anxiety in patients before elective general surgery. *International Journal of Pharmaceutical Research.* 2018;10(3):136-42. [View at Publisher] [DOI] [Google Scholar]
- Ghimire R, Poudel P. Preoperative anxiety and its determinants among patients scheduled for major surgery: a hospital based study. *J Anesthesiol.* 2018;6(2):57-60. [View at Publisher] [Google Scholar]
- Renaud-Roy E, Stöckle P-A, Maximos S, Brulotte V, Sideris L, Dubé P, et al. Correlation between incremental remifentanyl doses and the Nociception Level (NOL) index response after intraoperative noxious stimuli. *Can J Anesth.* 2019;66(9):1049-61. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Shirzad M, Nasiri E, Hesamirostami MH, Akbari H. Comparing the effects of rose essential oil and Benson relaxation technique on preoperative anxiety and hemodynamic status and postoperative complications in rhinoplasty candidates. *Complementary Medicine Journal.* 2021;11(2):180-91. [View at Publisher] [DOI] [Google Scholar]
- Vatankhah M, Melekshoar M. Comparison of the prophylactic effect of ibuprofen and intravenous ketorolac in pain control after upper extremity surgery: A Double-Blind Randomized Clinical Trial Study. *medical journal of mashhad university of medical sciences.* 2021;64(2):2692-705. [View at Publisher] [DOI] [Google Scholar]
- Lin CL, Hwang SL, Jiang P, Hsiung NH. Effect of music therapy on pain after orthopedic surgery-a systematic review and meta-analysis. *Pain Practice.* 2020;20(4):422-36. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Babatabar Darzi H, Vahedian-Azimi A, Ghasemi S, Ebadi A, Sathyapalan T, Sahebkar A. The effect of aromatherapy with rose and lavender on anxiety, surgical site pain, and extubation time after open-heart surgery: A double-center randomized controlled trial. *Phytother Res.* 2020;34(10):2675-84. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Grossman LD, Roscoe R, Shack AR, Committee DCCPGE. Complementary and alternative medicine for diabetes. *Can J Diabetes.* 2018;42(Suppl1):S154-61. [View at Publisher] [DOI] [PMID] [Google Scholar]
- Farzaneh M, Abbasijahromi A, Saadatmand V, Parandavar N, Dowlatkhan HR, Bahmanjahromi A. Comparative effect of nature-based sounds