

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/ajur

Original Article

Outcomes of surgical correction of Peyronie's disease with plaque excision and grafting: Comparison of testicular tunica vaginalis graft versus bovine pericardium graft

Ali Eslahi ^{a,b}, Faisal Ahmed ^{c,*}, Mohammad Reza Askarpour ^a,
 Hossein-Ali Nikbakht ^d, Iman Shamohammadi ^a,
 Payam Ghasemi ^a, Hanieh Alimardani ^a, Bahareh Ebrahimi ^b

^a Department of Urology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

^b Shiraz Geriatric Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

^c Urology Research Center, Al-Thora General Hospital, Department of Urology, School of Medicine, Ibb University of Medical Sciences, Ibb, Yemen

^d Social Determinants of Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

Received 5 October 2022; accepted 29 March 2023

Available online 2 August 2023

KEYWORDS

Peyronie's disease;
 Plaque excision;
 Testicular tunica vaginalis graft;
 Bovine pericardium graft

Abstract *Objective:* Peyronie's disease (PD) is an abnormal wound healing in the penile tunica albuginea. After fibrotic plaque excision, different graft materials have been used to repair the defects, but the optimal graft remains unknown. This study aimed to compare the functional outcomes of testicular tunica vaginalis grafts and bovine pericardium grafts in patients with severe PD.

Methods: A retrospective comparative study was conducted on 33 PD patients undergoing partial plaque excision and grafting from September 2015 to May 2021. The patients were divided into two groups depending on the type of graft used. For 15 patients in Group B, testicular tunica vaginalis grafts were used to repair the defect, while for 18 patients in Group A, bovine pericardium grafts were used. Data of the patient's age, comorbidities, sexual function, penile curvature, postoperative complications, need for further treatment, change in penile length, and satisfaction were gathered and compared between the groups. Sexual function was evaluated using the 5-item version of the International Index of Erectile Function (IIEF-5), and a

* Corresponding author.

E-mail address: fmaaa2006@yahoo.com (F. Ahmed).

Peer review under responsibility of Tongji University.

<https://doi.org/10.1016/j.ajur.2023.03.005>

2214-3882/© 2024 Editorial Office of Asian Journal of Urology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

hypoesthesia, orchialgia, and orchitis were reported in 17.3%, 56.5%, 8.3%, and 8.3%, respectively. The reported complication of bovine pericardium graft was cyst formation, graft contraction, infection, loss of sensation, and rarely urethral cutaneous fistula [12,19,24]. Conversely, in our study, the reported complications were hypoesthesia in three cases in each group, and infection, which was presented in one patient in the bovine pericardium group and managed with medical therapy. As expected, since all surgeries in our study were carried out by a single academic urologist with many years of experience in this field, we had lower rates of complications in comparison to other researchers.

In our study, the difference between IIEF-5 scores in the pericardial graft and testicular tunica vaginalis groups was statistically insignificant. In contrast, the mean baseline IIEF-5 scores in both groups were slightly decreased, which was statistically insignificant ($p=0.492$). Reviewing previous literature, we found disparities in the reported IIEF-5 scores after the surgery. For instance, Flores and his team [25] reported a 6-point reduction in IIEF-5 after the surgery. Similarly, Chung et al. [26] noted a decrease of 5 points in the IIEF-5 scores after PD graft surgery. On the contrary, Ainayev et al. [20] reported a 2-point improvement in the IIEF-5 score within 2 years after the surgery (mean 18.4 [SD 2.5] to 20.6 [SD 2.60]). Likewise, Liu et al. [8] reported improvement in the IIEF-5 score with the follow-up from 12 months to 43 months. Differences between the studies in factors like case selection, surgeon experience, patient orientation, and level of education could explain these disparities in the IIEF-5 scores.

Long-term studies highlight the risks of decreased erectile function in response to the progression of arteriogenic or veno-occlusive dysfunction. For example, Kalsi et al. [27] reported a 22% decreased erectile function within 5 years of follow-up. Taylor and Levine [28] reported a 21% postoperative ED in the follow-up. Additionally, a recent literature review regarding plaque incisions and grafting showed that 4.6%–67.4% of patients required medicine to have erections postoperatively, and 0%–11.8% were utterly unable to achieve erections [29]. On the other hand, current studies suggested penile prosthesis implantation with or without additional straightening maneuvers for severe PD and concomitant ED [6,30].

In our study and within the follow-up period, there were 11 (33.3%) patients (5 vs. 6 patients in groups A and B, respectively) who used PDE5-Is to attempt penetrative sex. The need for PDE5-Is may not reflect ED as a result of the grafting procedure but may also be related to age-related changes occurring over the mean follow-up period [15].

Regarding satisfaction with the results, most patients and their partners were satisfied in both groups without a statistically significant difference (77.8% in pericardium graft vs. 86.7% in tunica vaginalis graft). The overall satisfaction is in accordance with a recent review by Ragheb et al. [5], which reported an overall satisfaction rate of 91.2% in pericardium grafted and 100.0% in tunica vaginalis grafted patients. Another multicenter study also described long-term outcomes for grafting 157 patients with the bovine pericardium, with a 97% satisfaction rate among patients within 2 years after surgery [31]. Case selection might be the key to our research's highly successful outcomes and satisfaction

rates. Only those patients who met our criteria underwent surgery. Less strict exclusion criteria might lead to poorer outcomes which should be investigated in future studies. Additionally, we had an extended follow-up period compared to the previous studies, and having a single academic urologist during the follow-up visits helped us monitor and assess our patient's progress carefully while keeping patients motivated and involved in their treatments.

Despite including all PD patients who required graft surgery for 6 years between 2015 and 2021, the small number of cases in this study was our main limitation. We suggest further studies with a more extensive study population and better randomization.

5. Conclusion

Our results indicate that partial plaque excision and corporoplasty with testicular tunica vaginalis graft or bovine pericardium graft are equally effective treatments for men with clinically significant PD.

Author contributions

Study concept and design: Ali Eslahi.

Data acquisition: Payam Ghasemi, Hanieh Alimardani, Bahareh Ebrahimi.

Data analysis: Hossein-Ali Nikbakht, Iman Shamohammadi.

Drafting of the manuscript: Ali Eslahi, Faisal Ahmed.

Critical revision of the manuscript: Faisal Ahmed, Mohammad Reza Askarpour.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgments

The authors would like to thank the Student Research Committee of Shiraz University of Medical Sciences for financially supporting this research. They would also like to thank the Center for Development of Clinical Research of Nemazee Hospital and Dr. Nasrin Shokrpour for editorial assistance in language editing.

References

- [1] El-Sakka AI, Salabas E, Dinçer M, Kadioglu A. The pathophysiology of Peyronie's disease. *Arab J Urol* 2013;11:272–7.
- [2] Ostrowski KA, Gannon JR, Walsh TJ. A review of the epidemiology and treatment of Peyronie's disease. *Res Rep Urol* 2016;8:61–70.
- [3] Chen R, McCraw C, Lewis R. Plication procedures—excisional and incisional corporoplasty and imbrication for Peyronie's disease. *Transl Androl Urol* 2016;5:318–33.
- [4] Fiorillo A, Droghetti M, Palmisano F, Bianchi L, Piazza P, Sadini P, et al. Long-term outcomes after plaque incision and grafting for Peyronie's disease: comparison of porcine dermal and bovine pericardium grafts. *Andrology* 2021;9:269–76.
- [5] Ragheb A, Eraky A, Osmonov D. A decade of grafting techniques as a sole treatment for Peyronie's disease. *Andrology* 2020;8:1651–9.